arXiv:1206.3364v2 [nlin.SI] 28 Jun 2012

Ground-state analysis for a coupled-spin analogue of the $p + ip$-pairing Hamiltonian

Eduardo Mattei and Jon Links
Centre for Mathematical Physics
School of Mathematics and Physics
The University of Queensland, 4072
Australia

Abstract

We introduce a Hamiltonian for two interacting $su(2)$ spins. This model is an analogue of the Bethe ansatz solvable $p+ip$-pairing Hamiltonian in that it admits a duality, from which the Read-Green and Moore-Read lines are present in the model's phase diagram. We use a mean-field analysis and exact Bethe ansatz results to investigate the ground-state properties of the system in the thermodynamic limit, defined here as the limit of infinite spin (or highest weight). We also numerically study the energy gap, fidelity and entanglement. We find that certain ground-state properties in the thermodynamic limit are ill-defined due to issues concerning non-commuting limits of the system's coupling parameters and the thermodynamic limit. We comment on possible implications for the $p + ip$-pairing Hamiltonian.

1 Introduction

The $p+ip$-pairing Hamiltonian has come to some prominence through the work of Read and Green [1] who, through a mean-field analysis, exposed a change in the topological properties of the ground-state wavefunction of the Hamiltonian as a function of the coupling parameter. An exact Bethe ansatz solution of the Hamiltonian was provided in [2], and shortly after two detailed analyses were conducted in [3, 4]. However their conclusions were different, with [3] arguing that there are three ground-state phases of the system while [4] maintained that there are two. In both cases the author's singled out the *Moore-Read* and *Read-Green* lines of the system, which will be defined later, as displaying some unique properties. In [3] it was argued that both lines should be identified with quantum phase boundaries, whereas [4] claimed that only the Read-Green line represented a quantum phase boundary.

We consider a simplified Hamiltonian for two interacting $su(2)$ spins, which we present as an analogue of the $p + ip$-pairing Hamiltonian in that it displays both the Moore-Read and Read-Green lines in the phase diagram as a result of a duality relation. We then conduct an analysis of the system using a mean-field approximation and the exact Bethe ansatz solution to investigate ground-state properties in the thermodynamic limit, defined here as the limit of infinite spin (or highest weight). An advantage of dealing with this Hamiltonian of two interacting $su(2)$ spins is that analytic results are accessible. We also numerically solve the system to study the ground-state energy gap, fidelity [5, 6], and entanglement [7, 8], which have been used as indicators of quantum phase transitions in other simple bipartite quantum systems e.g. [9–11]. We find that certain ground-state properties in the thermodynamic limit are ill-defined, due to issues concerning non-commuting limits of the system's coupling parameters and the thermodynamic limit. We comment on some possible implications for the $p + ip$-pairing Hamiltonian.

2 The Hamiltonian

The $su(2)$ algebra has commutation relations

$$\left[S^+,\, S^-\right] = 2S^z, \qquad \left[S^z,\, S^\pm\right] = S^\pm. \tag{1}$$

Given a finite-dimensional module with highest weight s the action of the generators on the weight basis is

$$\begin{aligned}
S^z|s,m\rangle &= m|s,m\rangle, \\
S^+|s,m\rangle &= \sqrt{(s-m)(s+m+1)}|s,m+1\rangle, \\
S^-|s,m\rangle &= \sqrt{(s+m)(s-m+1)}|s,m-1\rangle
\end{aligned}$$

with $m = -s, -s+1, ..., s-1, s$.

We will study the following coupled-spin Hamiltonian

$$H = (1+G)(z_1^2(S_1^z+s) + z_2^2(S_2^z+s)) - G(z_1S_1^+ + z_2S_2^+)(z_1S_1^- + z_2S_2^-) \tag{2}$$

which acts on the $(2s+1)^2$-dimensional tensor product space with basis

$$\{|s,m_1\rangle \otimes |s,m_2\rangle \,:\, m_j = -s, -s+1, ..., s-1, s;\; j = 1,2\}. \tag{3}$$

Throughout we assume $z_1 \geq z_2 \geq 0$. For given s the model has three coupling parameters. One of these can be fixed by choosing the overall energy scale, e.g. setting $z_1 = 1$, which leaves two independent parameters z_2 and G. Define

$$\begin{aligned}
H_0 &= z_1^2(S_1^z+s_1) + z_2^2(S_2^z+s_2) \\
Q^+ &= z_1S_1^+ + z_2S_2^+ \\
Q^- &= z_1S_1^- + z_2S_2^- \\
B^+ &= \frac{1}{z_1}S_1^+ + \frac{1}{z_2}S_2^+ \\
S^z &= S_1^z + S_2^z
\end{aligned}$$

such that

$$H = (1+G)H_0 - GQ^+Q^- \tag{4}$$

and $[S^z,\, H] = 0$. From the commutation relations (1) the following equation can be shown to hold

$$\left[H,\, (B^+)^M\right] = MQ^+(B^+)^{M-1}(1+GM+2GS^z).$$

If a state $|\psi\rangle$ has spin m (the eigenvalue of S^z) the above means that choosing $1+GM+2Gm = 0$, so $M = -2m-1/G$, the two states $|\psi\rangle$ and $|\tilde{\psi}\rangle = (B^+)^M|\psi\rangle$ have the same energy. We call the states $|\psi\rangle$ and $|\tilde{\psi}\rangle$ *dual* states. Since the state $|\psi\rangle$ has spin m the state $|\tilde{\psi}\rangle$ has spin $m+M = -m-1/G$. We identify the following cases:

- The dual of the lowest weight state $|\psi\rangle = |s,-s\rangle \otimes |s,-s\rangle$ occurs when $M = 4s-1/G$ leading to $|\tilde{\psi}\rangle$ having spin $m = 2s-1/G$. This is referred to as the Moore-Read line, on which the ground-state energy is zero.
- If $M = 0$ then a state is self-dual when $m = -1/(2G)$. This is referred to as the Read-Green line.

Note that the above definitions for the Moore-Read and Read-Green lines are *independent* of the parameters z_1 and z_2, analogous to the case of the $p+ip$-pairing Hamiltonian where these lines are independent of the single particle momentum distribution [2–4]. The Hamiltonian (2) may be viewed as a degenerate, two-level limit of the $p + ip$-pairing Hamiltonian on a restricted Hilbert space, which has been studied in [12].

Our main objective is to investigate ground-state properties of the model in the *thermodynamic limit* which we define by a sequence of ordered triples (G, m, s) such that $G \to 0$, $m \to \infty$, $s \to \infty$ with $\mu = m/(2s)$, $g = 4sG$ held constant. In terms of these parameters the Moore-Read line is given by $\mu = 1 - 2/g$ while the Read-Green line is $\mu = -1/g$.

3 Mean-field analysis

The initial study by Read and Green [1] on the $p + ip$-pairing Hamiltonian was undertaken using a mean-field analysis. Here we start by taking an analogous study for the coupled-spin model. The Hamiltonian (2) can be approximated by a Hamiltonian with linear terms through

$$AB \approx A\langle B\rangle + \langle A\rangle B - \langle A\rangle\langle B\rangle$$

where $\langle A\rangle$ denotes the expectation value of the operator A. The mean-field approximation which linearises the Hamiltonian (2) is

$$\begin{aligned} H_{\mathrm{mf}} &= (1+G)s(z_1^2+z_2^2) + \rho m + ((1+G)z_1^2-\rho)S_1^z + ((1+G)z_2^2-\rho)S_2^z \\ &\quad - \frac{\hat{\Delta}}{2}(z_1S_1^+ + z_2S_2^+) - \frac{\hat{\Delta}^*}{2}(z_1S_1^- + z_2S_2^-) + \frac{\Delta^2}{4G}, \end{aligned}$$

where ρ is a Lagrange multiplier (the analogue of the chemical potential) which is introduced as the mean-field approximation breaks the conservation of S^z, and

$$\hat{\Delta} = 2G\langle z_1S_1^- + z_2S_2^-\rangle.$$

Setting $\Delta = |\hat{\Delta}|$ the ground state energy is found to be

$$E_0 = (1+G)s(z_1^2+z_2^2) + \rho m + \frac{\Delta^2}{4G} - s\sqrt{((1+G)z_1^2-\rho)^2 + z_1^2\Delta^2} - s\sqrt{((1+G)z_2^2-\rho)^2 + z_2^2\Delta^2}.$$

Using the Hellmann-Feynman theorem we can determine the mean-field expectation values:

$$\begin{aligned} &\left\langle \frac{\partial H_{\mathrm{mf}}}{\partial G} \right\rangle = \frac{\partial E_0}{\partial G} \\ \Longrightarrow \qquad & z_1^2\langle S_1^z\rangle + z_2^2\langle S_2^z\rangle + \frac{\Delta^2}{2G^2} = sz_1^2\frac{(1+G)z_1^2 - \rho + \Delta^2/G}{\sqrt{((1+G)z_1^2-\rho)^2 + z_1^2\Delta^2}} + sz_2^2\frac{(1+G)z_2^2 - \rho + \Delta^2/G}{\sqrt{((1+G)z_2^2-\rho)^2 + z_2^2\Delta^2}}. \end{aligned}$$

Taking the thermodynamic limit gives

$$\frac{2}{g} = \frac{z_1^2}{\sqrt{(z_1^2-\rho)^2 + z_1^2\Delta^2}} + \frac{z_2^2}{\sqrt{(z_2^2-\rho)^2 + z_2^2\Delta^2}}. \tag{5}$$

Also

$$\begin{aligned} &\left\langle \frac{\partial H_{\mathrm{mf}}}{\partial \rho} \right\rangle = \frac{\partial E_0}{\partial \rho} \\ \Longrightarrow \qquad & -\langle S_1^z\rangle - \langle S_2^z\rangle = \frac{s((1+G)z_1^2-\rho)}{\sqrt{((1+G)z_1^2-\rho)^2 + z_1^2\Delta^2}} + \frac{s((1+G)z_2^2-\rho)}{\sqrt{((1+G)z_2^2-\rho)^2 + z_2^2\Delta^2}}. \end{aligned}$$

Taking the thermodynamic limit and using (5) gives

$$\frac{2}{g} + 2\mu = \frac{\rho}{\sqrt{(z_1^2-\rho)^2 + z_1^2\Delta^2}} + \frac{\rho}{\sqrt{(z_2^2-\rho)^2 + z_2^2\Delta^2}}. \tag{6}$$

Returning to the energy expression in the limit $s \to \infty$, and taking into account Eqs. (5,6), we obtain the intensive energy expression

$$\begin{aligned}
e_0 &= \lim_{s\to\infty} \frac{E_0}{s} \\
&= (z_1^2 + z_2^2) + 2\rho\mu + \frac{\Delta^2}{g} - \sqrt{(z_1^2-\rho)^2 + z_1^2\Delta^2} - \sqrt{(z_2^2-\rho)^2 + z_2^2\Delta^2} \\
&= z_1^2\left(1 - \frac{1}{2}\frac{2z_1^2 + \Delta^2 - 2\rho}{\sqrt{(z_1^2-\rho)^2 + z_1^2\Delta^2}}\right) + z_2^2\left(1 - \frac{1}{2}\frac{2z_2^2 + \Delta^2 - 2\rho}{\sqrt{(z_2^2-\rho)^2 + z_2^2\Delta^2}}\right). \qquad (7)
\end{aligned}$$

It may be verified that setting $\rho = \Delta^2/4$ in (5,6) and combining these equations leads to the equation of the Moore-Read line. Furthermore, setting $\rho = \Delta^2/4$ in (7) gives $e_0 = 0$ in agreement with the result that ground states on the Moore-Read line are dual to the lowest weight state.

Recalling the assumption $z_1 > z_2$ the mimimum excitation energy (i.e. the gap) is

$$\Delta E = \sqrt{(z_2^2-\rho)^2 + z_2^2\Delta^2}$$

which vanishes when $z_2 = \rho = 0$. However turning to (6) we see that the limits $z_2 \to 0$ and $\rho \to 0$ do not commute in that equation. First setting $\rho = 0$ leads to the condition of the Read-Green line. Alternatively first setting $z_2 = 0$ yields

$$\mu = \frac{1}{2}\lim_{\rho\to 0}\frac{\rho}{|\rho|} - \frac{1}{g} \qquad (8)$$

which shows a discontinuity as ρ passes through zero. This indicates that while gapless excitations occur in the limit $z_2, \rho \to 0$, where they are to be found in the phase diagram will depend on how these two limits are taken (at least within the mean-field approximation).

In the analogous study of Read and Green [1], a similar type of discontinuity was also observed in a particular number operator expectation value. Through this observation the Read-Green line was identified with a topological phase transition. For the simplified coupled-spin model we will next conduct an analysis using exact results. In this manner we will highlight some subtle aspects regarding the ground-state properties of the system, which appear to be beyond the scope of a mean-field analysis.

4 Exact results

4.1 The limiting case $z_2 = 0$

Here we set $z_2 = 0$ from the outset, and fix the energy scale by letting $z_1 = 1$, in which case the Hamiltonian is simply

$$H = (1+G)(S_1^z + s) - GS_1^+S_1^-.$$

The above Hamiltonian is diagonal in the basis (3) with eigenvalues

$$E = m_1 + s - G(s^2 - m_1^2). \qquad (9)$$

For each value of $m = m_1 + m_2$ we can determine the ground state energy, and the gap, by considering where level crossings occur. We need to consider several different cases. Starting with $m = m_1 + m_2 < 0$ we have

$$\begin{aligned}
&m_1 = -s, && m_2 = m + s && \Longrightarrow && E = 0, \\
&m_1 = -s + 1, && m_2 = m + s - 1 && \Longrightarrow && E = (1+G) - 2sG.
\end{aligned}$$



Figure 1: Phase diagram obtained from duality. The solid line $\mu = 1 - 2/g$ is the Moore-Read line, on which the intensive ground-state energy is zero. The dotted line $\mu = -1/g$ is the Read-Green line. The mean-field analysis predicts that there are gapless excitations on the Read-Green line as $z_2 \to 0$, but also that the limits $\mu \to -1/g$ and $z_2 \to 0$ do not commute. With respect to the duality discussions of Sect. 2, the Moore-Read line denotes states dual to the lowest weight state ($\mu = -1$), while the Read-Green line denotes states which are self-dual. Note that neither boundary line depends on the variables z_1 and z_2.

In the thermodynamic limit it is seen that that for $m < 0$ a first level crossing occurs at

$$g = 2. \tag{10}$$

Next consider

$$\begin{aligned} m_1 &= s + m - 1, \qquad m_2 = -s + 1 \qquad \Longrightarrow \qquad E = 2s + m - 1 + G((m-1)^2 + 2s(m-1)),\\ m_1 &= s + m, \qquad m_2 = -s \qquad \Longrightarrow \qquad E = 2s + m + G(m^2 + 2sm). \end{aligned}$$

Here there is an energy gap

$$\Delta E = |1 + G(2m - 1 + 2s)|.$$

In the thermodynamic limit the final level crossing occurs when

$$\mu = -\frac{1}{2} - \frac{1}{g} \tag{11}$$

which requires that $\mu < -1/2$.

Now we consider the case for $m \geq 0$:

$$\begin{aligned} m_1 &= -s + m, \qquad m_2 = s \qquad \Longrightarrow \qquad E = m + G(m^2 - 2sm),\\ m_1 &= -s + m + 1, \qquad m_2 = s - 1 \qquad \Longrightarrow \qquad E = m + 1 + G((m+1)^2 - 2s(m+1)). \end{aligned}$$

The gap

$$\Delta E = |1 + G(2m + 1 - 2s)|$$

goes to zero in the thermodynamic limit when

$$\mu = \frac{1}{2} - \frac{1}{g}. \tag{12}$$

This requires that $0 \leq \mu < 1/2$. In the region bounded by the equations (10,11,12) the level crossings become dense in the thermodynamic limit, giving rise to a gapless phase. The following relations hold for the gap between the ground and first-excited states in the regions depicted in the phase diagram of Fig. 2:

$$\begin{aligned} \mathrm{I}: &\quad \Delta E = 1 + g(\mu - 1/2), && (13)\\ \mathrm{II}: &\quad \Delta E = 1 - g/2, && (14)\\ \mathrm{III}: &\quad \Delta E = -1 - g(\mu + 1/2), && (15)\\ \mathrm{IV}: &\quad \Delta E = 0. && (16) \end{aligned}$$



Figure 2: Ground-state phase diagram for $z_2 = 0$. Expressions for the gap between the ground and first-excited states in the thermodynamic limit are given by Eqs. (13-16). Region IV is gapless.

There is clear discrepancy between Figs. 1 and 2, in particular for the predicted gapless regions. In deriving the two phase diagrams we consider behaviours as $s \to \infty$, $m \to \infty$, $G \to 0$ with $g = 4sG$, $\mu = m/(2s)$, and as $z_2 \to 0$, yielding phase boundary lines as functions of the coupling parameter g. So far there are indications that the limiting behaviours do not always commute. Similar issues with non-commuting limits have been observed in the different context of attractive bosons [13]. In the following three Subsections we will appeal to further exact results to gain a better insight into the properties of the system.

4.2 The case $m = 1 - 2s$

In the sector where $m = 1 - 2s$ the Hilbert space is two-dimensional and exact results are easily extracted. In matrix form, with respect to the basis

$$\{|s, -s+1\rangle \otimes |s, -s\rangle,\ |s, -s\rangle \otimes |s, -s+1\rangle\},$$

the Hamiltonian is

$$H = \begin{pmatrix} (1 + G - 2sG)z_1^2 & -2sGz_1z_2 \\ -2sGz_1z_2 & (1 + G - 2sG)z_2^2 \end{pmatrix} \tag{17}$$

with the ground-state energy

$$E_{[m=1-2s]} = \frac{(1 + g/(4s) - g/2)(z_1^2 + z_2^2)}{2} - \frac{1}{2}\sqrt{(1 + g/(4s) - g/2)^2(z_1^2 - z_2^2)^2 + g^2 z_1^2 z_2^2} \tag{18}$$

and energy gap to the excited state

$$\Delta E = \sqrt{(1+g/(4s)-g/2)^2(z_1^2-z_2^2)^2+g^2z_1^2z_2^2}.$$

Here $\mu \to -1$ in the thermodynamic limit, which at $g = 1$ corresponds to the limit where the Moore-Read and Read-Green lines meet. It is found that

$$\lim_{z_2\to 0}\lim_{g\to 2}\lim_{s\to\infty}\Delta E = 0$$

with the result being invariant with respect to interchanging the order of the limits. Moreover, as $s \to \infty$ the *only* solution for $\Delta E = 0$ when $z_1 \neq 0$ is given by $z_2 = 0$, $g = 2$, meaning that gapless excitations *do not* occur at the Read-Green line in this limit $\mu \to -1$. The behaviour accords with Fig. 2 rather than Fig. 1.

Next consider

$$\frac{\partial(\Delta E)}{\partial g} = \frac{g(z_1^2+z_2^2)^2-(2-(1-g+g/(4s))/s)(z_1^2-z_2^2)^2}{4\Delta E}.$$

We find

$$\lim_{z_2\to 0}\lim_{g\to 2}\lim_{s\to\infty}\frac{\partial(\Delta E)}{\partial g} = 0, \qquad \lim_{z_2\to 0}\lim_{s\to\infty}\lim_{g\to 2}\frac{\partial(\Delta E)}{\partial g} = 0,$$
$$\lim_{g\to 2^+}\lim_{z_2\to 0}\lim_{s\to\infty}\frac{\partial(\Delta E)}{\partial g} = \frac{z_1^2}{2}, \qquad \lim_{g\to 2^+}\lim_{s\to\infty}\lim_{z_2\to 0}\frac{\partial(\Delta E)}{\partial g} = \frac{z_1^2}{2},$$
$$\lim_{g\to 2^-}\lim_{z_2\to 0}\lim_{s\to\infty}\frac{\partial(\Delta E)}{\partial g} = -\frac{z_1^2}{2}, \qquad \lim_{g\to 2^-}\lim_{s\to\infty}\lim_{z_2\to 0}\frac{\partial(\Delta E)}{\partial g} = -\frac{z_1^2}{2},$$
$$\lim_{s\to\infty}\lim_{z_2\to 0}\lim_{g\to 2}\frac{\partial(\Delta E)}{\partial g} = -\frac{z_1^2}{2}, \qquad \lim_{s\to\infty}\lim_{g\to 2}\lim_{z_2\to 0}\frac{\partial(\Delta E)}{\partial g} = -\frac{z_1^2}{2}.$$

This establishes that the non-commutativity of limits is not just an artefact of the mean-field approximation, but is an intrinsic property of the model.

4.3 Non-analytic behaviour on the Moore-Read line

Next we will use the duality properties of the model to show that non-analytic behaviour is found on the Moore-Read line appearing in Fig. 1. Our approach is through a study of how the intensive ground-state energy $e_0(g)$ varies with g while z_1, z_2 and μ are held constant.

First we consider two ground states of the system at different values of the coupling parameter:

$$G^{(1)} = \frac{1}{2s-m}, \qquad G^{(2)} = \frac{1}{2s-m-1}$$

such that $\Delta G = G^{(2)} - G^{(1)} = G^{(1)}G^{(2)}$. We also have $\Delta g = 4s(G^{(2)} - G^{(1)}) = 4sG^{(1)}G^{(2)}$. It is found that in the thermodynamic limit

$$\lim_{s\to\infty} g^{(1)} = \lim_{s\to\infty} 4sG^{(1)} = \lim_{s\to\infty}\frac{4s}{2s-m} = \lim_{s\to\infty}\frac{2}{1-\mu} = \frac{2}{1-\mu},$$
$$\lim_{s\to\infty} g^{(2)} = \lim_{s\to\infty} 4sG^{(2)} = \lim_{s\to\infty}\frac{4s}{2s-m-1} = \lim_{s\to\infty}\frac{2}{1-\mu-1/(2s)} = \frac{2}{1-\mu}$$

so we hereafter set

$$g_{\mathrm{MR}} = \frac{2}{1-\mu}.$$

Then

$$\lim_{s\to\infty} 4s\Delta g = g_{\mathrm{MR}}^2. \tag{19}$$

The ground state corresponding to $g^{(1)} = 4sG^{(1)}$ is on the Moore-Read line, and so has energy $E_0(g^{(1)}) = 0$. The ground state corresponding to $g^{(2)} = 4sG^{(2)}$ is dual to the ground state of (17), and so has energy $E_0(g^{(2)}) = E_{[m=1-2s]}$ as given by (18). We also have

$$e_0(g_{\rm MR}) = \lim_{s\to\infty} \frac{E_0(g^{(1)})}{s} = \lim_{s\to\infty} \frac{E_0(g^{(2)})}{s} = 0.$$

Now consider in general terms

$$\begin{aligned} \frac{\partial e_0(g)}{\partial g} &= \lim_{\delta\to 0} \frac{e_0(g+\delta) - e_0(g)}{\delta} \\ &= \lim_{\delta\to 0} \lim_{s\to\infty} \frac{E_0(g+\delta) - E_0(g)}{s\delta}. \end{aligned}$$

One could alternatively define

$$\begin{aligned} \frac{\partial e_0(g)}{\partial g} &= \lim_{s\to\infty} \frac{1}{s} \frac{\partial E_0(g)}{\partial g} \\ &= \lim_{s\to\infty} \lim_{\delta\to 0} \frac{E_0(g+\Delta g) - E_0(g)}{s\delta}. \end{aligned}$$

The definitions will be equivalent if the limits $\delta \to 0$ and $s \to \infty$ commute.

If the limits $g \to g_{\rm MR}$ and $s \to \infty$ commute we have

$$\begin{aligned} \frac{\partial e_0(g_{\rm MR})}{\partial g} &= \lim_{\delta\to 0} \frac{e_0(g_{\rm MR}+\delta) - e_0(g_{\rm MR})}{\delta} \\ &= \lim_{\delta\to 0} \frac{e_0(g_{\rm MR}+\delta)}{\delta} \\ &= \lim_{\delta\to 0} \lim_{s\to\infty} \frac{E_0(g^{(1)}+\delta)}{s\delta} \\ &= \lim_{s\to\infty} \frac{E_0(g^{(1)}+\Delta g)}{s\Delta g} \\ &= \lim_{s\to\infty} \frac{1}{s\Delta g} \lim_{s\to\infty} E_0(g^{(2)}) \\ &= \frac{(2-g_{\rm MR})(z_1^2+z_2^2)}{g_{\rm MR}^2} - \frac{2}{g_{\rm MR}^2}\sqrt{(1-g_{\rm MR}/2)^2(z_1^2-z_2^2)^2 + g_{\rm MR}^2 z_1^2 z_2^2} \end{aligned}$$

where in the last step we use (18,22). Alternatively

$$\begin{aligned} \frac{\partial e_0(g_{\rm MR})}{\partial g} &= \lim_{\delta\to 0} \frac{e_0(g_{\rm MR}+\delta) - e_0(g_{\rm MR})}{\delta} \\ &= \lim_{\delta\to 0} \frac{e_0(g_{\rm MR}+\delta)}{\delta} \\ &= \lim_{\delta\to 0} \lim_{s\to\infty} \frac{E_0(g^{(2)}+\delta)}{s\delta} \\ &= -\lim_{s\to\infty} \frac{E_0(g^{(2)}-\Delta g)}{s\Delta g} \\ &= -\lim_{s\to\infty} \frac{1}{s\Delta g} \lim_{s\to\infty} E_0(g^{(1)}) \\ &= 0. \end{aligned}$$

So assuming that the limits $s \to \infty$ and $g \to g_{\rm MR}$ commute leads to the partial derivative being discontimuous.

4.4 Bethe ansatz solution

As in the case of the $p + ip$ model [2], the Hamiltonian (2) admits a Bethe ansatz solution. The energy spectrum is given by

$$E = (1 + G) \sum_{k=1}^{M} y_k$$

where the y_k, $k = 1, ..., M$, are roots of the Bethe ansatz equations

$$\frac{G^{-1} + 2M - 4s - 1}{y_k} + \frac{2s}{y_k - z_1^2} + \frac{2s}{y_k - z_2^2} = \sum_{j \neq k}^{M} \frac{2}{y_k - y_j} \tag{20}$$

and the total spin is $m = M - 2s$. For each solution of the Bethe ansatz equations the associated eigenstate is

$$|\psi\rangle = \prod_{k=1}^{M} C(y_k) \left(|s, -s\rangle \otimes |s, -s\rangle\right)$$

where

$$C(y) = \frac{z_1 S_1^+}{y - z_1^2} + \frac{z_2 S_2^+}{y - z_2^2}. \tag{21}$$

The validity of this solution can be verfied in a direct manner, using only the commutation relations (1). We omit the details.

Following [3, 4], we look to solve the Bethe ansatz equations in the thermodynamic limit under the assumption that in this limit the roots of (20) are densely distributed on a curve Γ in the complex plane which is invariant under reflections about the real axis. Introducing $r(y)$ as the density function on Γ for the roots, we divide (20) by $2s$ and take the limit $s \to \infty$ to obtain

$$\frac{2g^{-1} + 2\mu}{y} + \frac{1}{y - z_1^2} + \frac{1}{y - z_2^2} = 2P \int_\Gamma |dy'| \frac{r(y')}{y - y'} \tag{22}$$

where P refers to the Cauchy principal value for the integral. To solve the problem we define an analytic function $h(y')$ outside Γ such that

$$r(y') = h_+(y') - h_-(y') \tag{23}$$

where $h_+(y')$ and $h_-(y')$ are the limiting values of $h(y')$ to the right and left of Γ. We can then rewrite (22) as

$$\frac{2g^{-1} + 2\mu}{y} + \frac{1}{y - z_1^2} + \frac{1}{y - z_2^2} = 2 \oint_L dy' \frac{h(y')}{y - y'} \tag{24}$$

where L encircles Γ. Next we take the following ansatz:

$$h(y) = \frac{R(y)}{2\pi i} \left[\frac{\phi(z_1)}{z_1^2 - y} + \frac{\phi(z_2)}{z_2^2 - y} + \frac{S}{y} \right] \tag{25}$$

where

$$R(y) = \sqrt{(y - a)(y - b)} \tag{26}$$

with a, $b = a^*$ denoting the end points of the arc Γ. Replacing (25) in (24) and evaluting the contour integral by residues leads to the conditions

$$\phi(z_j) = \frac{1}{2\sqrt{(z_j^2 - a)(z_j^2 - b)}},$$

$$S = \phi(z_1) + \phi(z_2),$$

$$S = \frac{g^{-1} + \mu}{\sqrt{ab}}.$$

These combine to give

$$\frac{2}{g} + 2\mu = \frac{\sqrt{ab}}{\sqrt{(z_1^2 - a)(z_1^2 - b)}} + \frac{\sqrt{ab}}{\sqrt{(z_2^2 - a)(z_2^2 - b)}}. \tag{27}$$

We may also compute

$$\begin{aligned}
\mu &= -1 + \int_\Gamma |dy'|\, r(y') \\
&= -1 + \oint_L dy'\, h(y') \\
&= \left(\frac{a+b}{2}\right) S - z_1^2 \phi(z_1) - z_2^2 \phi(z_2) - \frac{(\sqrt{a} - \sqrt{b})^2}{2} S \\
\Longrightarrow \qquad \frac{2}{g} &= \frac{z_1^2}{\sqrt{(z_1^2 - a)(z_1^2 - b)}} + \frac{z_2^2}{\sqrt{(z_2^2 - a)(z_2^2 - b)}}.
\end{aligned} \tag{28}$$

Identifying $\rho = \sqrt{ab}$ and $\Delta^2 = a + b + 2\sqrt{ab}$, Eqs. (27,28) are precisely the mean-field equations (6,5) respectively. This would suggest that the mean-field equations should be exact in the thermodynamic limit as they are reproducible from the exact Bethe ansatz solution. However, the above calculations assume that in the thermodynamic limit the roots of the Bethe ansatz equations become dense on a curve Γ. This assumption is not always valid. Suppose that the curve Γ closes such that the end points a, b become real and equal. Then Eqs. (27,28) reduce to

$$\frac{2}{g} + 2\mu = \frac{|a|}{|z_1^2 - a|} + \frac{|a|}{|z_2^2 - a|}, \tag{29}$$

$$\frac{2}{g} = \frac{z_1^2}{|z_1^2 - a|} + \frac{z_2^2}{|z_2^2 - a|}. \tag{30}$$

If $a \geq 0$ the only solution is $\mu = -1$, whereas if $a < 0$ we obtain the equation of the Moore-Read line $\mu = 1 - 2g^{-1}$. But the ground-state roots on the Moore-Read line are all zero, as the ground state is dual to the lowest weight state. Thus taking $s \to \infty$ after $g \to g_{\mathrm{MR}}$ leads to a delta function distribution of roots at the origin, while taking $g \to g_{\mathrm{MR}}^{-}$ after $s \to \infty$ leads to the roots being distributed on a closed curve enclosing the origin. The non-commutativity of these limits has previously been observed in the $p + ip$-pairing Hamiltonian. In [2,3] this was referred to as a zeroth order transition. We may identify the Moore-Read line as the boundary at which mean-field theory breaks down. In the region immediately beyond the Moore-Read line we cannot assume that the ground-state roots of the Bethe ansatz equations are densely distributed on a single curve, in order to produce the mean-field equations from the Bethe ansatz solution.

5 Indicators of quantum phase transitions

While there have been numerous studies of quantum phase transitions, there still lacks a definitive property which may be used to identify them. In the remaining Subsections we study various methods to gain an insight into the nature of possible quantum phase transitions for this system. Using numerical diagonalisation of the Hamiltonian (2), below we compute the energy gap, fidelity, and entanglement (as given by the von Neumann entropy). Each quantity is plotted as a function of the parameter g^{-1}. We fix $\mu = -0.99$, $z_1 = 1$, and adopt the parameterisation $z_2 = (2000s)^{-1/2}$ such that $z_2 \to 0$ as $s \to \infty$. We focus on this case as it permits us to compactly cover the regions II, III and IV in Fig. 2.

5.1 Energy Gap

In Fig. 3 we plot the energy gap. There are two different points of minima corresponding to the boundary lines between region III and IV, and between region IV and II, in Fig. 2. It is seen in the inset that as s increases the energy gap decreases in region IV of Fig. 2. The results appear more consistent with Fig. 2 rather than Fig. 1, as to be expected in view of the results from Subsect. 4.2.



Figure 3: Energy gap between the ground state and the first excited state as a function of g^{-1}, with $\mu = -0.99$. The results shown in the inset indicate the tendency towards vanishing gap in region IV as $s \to \infty$. The solid black line represents the analytic curve for the energy gap as given by Eqs. (14,15) for when $z_2 = 0$.

5.2 Fidelity

Another means to investigate quantum phase transitions is using fidelity [5,6], a concept widely used in quantum information theory. The fidelity of two states is simply defined as the modulus of the inner product

$$\mathcal{F}(\psi, \phi) = |\langle\psi|\phi\rangle|. \tag{31}$$

Computing this quantity for ground-state vectors associated with Hamiltonians differing only by a small change in the coupling parameter, the fidelity may be sensitive to variations of the coupling parameter. Generally a minimum in the fidelity points to a rapid change in the ground-state properties, which may indicate a critical point of the system.

Here we take two coupling values

$$(g^{\pm})^{-1} = g^{-1}(1 \pm \gamma)$$

with $\gamma = 0.001$ to compute the fidelity of the ground states associated with $g^{\pm}$, and then consider the behaviour of the fidelity as g is varied. As is seen in Fig. 4, for values of the parameter g^{-1} above 0.5 and values below 0.49, the fidelity is close to 1, but suffers an abrupt change at these two points. The two points correspond to the boundary lines between regions II and IV and regions IV and III for $\mu = -0.99$ in the phase diagram Fig. 2. Also, it can be seen that the minimum of the fidelity is decreasing as s increases.



Figure 4: Ground-state fidelity as a function of g^{-1}, with $\mu = -0.99$ and $\gamma = 0.001$ for different values of s as indicated in the legend. The results show significantly decreasing fidelity in region IV as $s \to \infty$.

5.3 Entanglement

The coupled-spin model is naturally seen as a bipartite system. In this case we can use the von Neumann entropy of the reduced density operator to quantify the entanglement of the system [7,9–11]. Using the fact that the total spin m is conserved, we can write a general state of the system (for $m \leq 0$) in terms of (3) by

$$|\Psi\rangle = \sum_{j=-s}^{m+s} d_j |s,j\rangle \otimes |s,m-j\rangle, \tag{32}$$

for some complex numbers d_j. The density operator for state (32) is given by

$$\rho = |\Psi\rangle\langle\Psi| = \sum_{i,j=-s}^{m+s} d_i d_j^* |s,i\rangle\langle s,j| \otimes |s,m-i\rangle\langle s,m-j|. \tag{33}$$

We take the partial trace with respect to the spin sector 2 yields the reduced density operator for the spin sector 1:

$$\rho_1 = \mathrm{tr}_2(\rho) = \sum_{j=-s}^{m+s} |d_j|^2 |s,j\rangle\langle s,j|. \tag{34}$$

The von Neumann entropy of entanglement for the state is given by

$$\mathcal{E}(\rho_1) = -\mathrm{tr}[\rho_1 \log_2(\rho_1)] = -\sum_{j=-s}^{m+s} |d_j|^2 \log_2(|d_j|^2). \tag{35}$$

Using the formula (35) we compute the entanglement of the ground state with $\mu = -0.99$ as function of g^{-1}, as shown in Fig. 5. We can identify two abrupt changes in the entanglement value about the points $g^{-1} = 0.49$ and $g^{-1} = 0.5$ in agreement with energy gap and fidelity analysis. Note that the entanglement variation gets sharper as the value of s is increased.



Figure 5: Ground-state entanglement as function of g^{-1} for $\mu = -0.99$ for different values of s as indicated in the legend. The results show significantly increasing entanglement in region IV as $s \to \infty$.

6 Conclusion

The mean-field analysis of the model predicted the ground-state phase diagram shown in Fig. 1 displaying the Moore-Read line, where the ground-state energy is zero, and Read-Green line, on which the gap goes to zero as $z_2 \to 0$. These lines also appear naturally with regards to the duality described in Sect. 2 and are *independent* of the variables z_1 and z_2. This is exactly the same phase diagram as that found for the $p + ip$-pairing Hamiltonian [2–4]. However starting with a finite-size system, setting $z_2 = 0$, and then taking the thermodynamic limit gave the ground-state phase diagram shown in Fig. 2. In Sect. 4 we turned to the exact Bethe ansatz solution for the model and through it found features associated with both phase diagrams. In the limit $\mu \to -1$ the gap was minimised at $g = 2$ as $z_2 \to 0$, in agreement with Fig. 2. Upon further analysis we also saw that the limits $s \to \infty$ and $z_2 \to 0$ do not always commute, with the derivative of the energy gap being ill-defined in this limit as $g \to 2$. On the other hand the Moore-Read line of Fig. 1 was found to exhibit non-analytic behaviour if the limits $s \to \infty$ and $g \to g_{\mathrm{MR}}$ were assumed to be commutative. Using the exact Bethe ansatz solution we argued that the Moore-Read line signifies the breakdown of the validity of mean-field solution. Finally we numerically studied the gap, fidelity, and entanglement to illustrate that signatures of the phase diagram in Fig. 2 were found in these quantities.

The complexity observed here in the simplified coupled-spin model highlights the need for deeper investigations into characterising the phase diagram of the $p + ip$-pairing Hamiltonian, in a manner which fully takes into account the peculiarities presented by the non-commuting limits. We expect that this is particularly the case in the so-called *weak-pairing* phase, which lies between the Moore-Read and Read-Green lines. It is in this region that the ground-state roots of the Bethe ansatz equations cannot be uniquely described by a density function, with support on a connected curve Γ, which varies smoothly with g. In this region there are limiting root distributions which have delta function terms at the origin, as a result of the duality, that are not connected to the distribution of the remaining roots.

Acknowledgements

EM is supported by CAPES (Brazil). JL is supported by the Australian Research Council through Discovery Project DP110101414. JL thanks Germán Sierra for numerous insightful discussions on the $p + ip$-pairing Hamiltonian.

References

[1] N. Read, D. Green, Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect, Phys. Rev. B **61** (2000) 10267.

[2] M. Ibañez, J. Links, G. Sierra, S.-Y. Zhao, Exactly solvable pairing model for superconductors with $p_x + ip_y$-wave symmetry, Phys. Rev. B **79** (2009) 180501.

[3] C. Dunning, M. Ibañez, J. Links, G. Sierra, S.-Y. Zhao, Exact solution of the $p + ip$ pairing Hamiltonian and a hierarchy of integrable models, J. Stat. Mech.: Theor. Exp. (2010) P08025.

[4] S.M.A. Rombouts, J. Dukelsky, G. Ortiz, Quantum phase diagram of the integrable $p_x + ip_y$ fermionic superfluid, Phys. Rev. B **82** (2010) 224510.

[5] P. Zanardi, N. Paunković, Ground state overlap and quantum phase transitions, Phys. Rev. E **74** (2006) 031123.

[6] H.-Q. Zhou, J.P. Barjaktarevič, Fidelity and quantum phase transitions, J. Phys. A: Math. Theor. **41** (2007) 412001.

[7] T.J. Osborne, M.A. Nielsen, Entanglement in a simple quantum phase transition, Phys. Rev. A **66** (2002) 032110.

[8] A. Osterloh, L. Amico, G. Falci, R. Fazio, Scaling of entanglement close to a quantum phase transition, Nature **416** (2002) 608.

[9] A.P. Hines, R.H. McKenzie, G.J. Milburn, Entanglement of two-mode Bose–Einstein condensates, Phys. Rev. A **67** (2003) 013609.

[10] F. Pan, J.P. Draayer, Quantum critical behavior of two coupled Bose–Einstein condensates, Phys. Lett. A **339** (2005) 403.

[11] D. Rubeni, A. Foerster, E. Mattei, I. Roditi, Quantum phase transitions in Bose–Einstein condensates from a Bethe ansatz perspective, Nucl. Phys. B **856** (2012) 698.

[12] I. Marquette, J. Links, Generalised Heine-Stieltjes and Van Vleck polynomials associated with degenerate, integrable BCS models, arXiv:1206.2988.

[13] N. Oelkers, J. Links, Ground-state properties of the attractive one-dimensional Bose-Hubbard model, Phys. Rev. B **75** (2007) 115119.